



MEDICAL ABORTION ON YOUR TERMS:

Good News for Patients

WHAT IS MEDICAL ABORTION?

Medical abortion, approved by the US Food and Drug Administration (FDA), is a proven method to end a pregnancy while it is still in its early stage — within 70 days from the first day of your last menstrual period. Used reliably in the US for more than 20 years by over five million people, medical abortion is private, safe, and non-invasive. It involves taking two types of pills: mifepristone, which blocks a hormone needed for pregnancy to continue, and misoprostol, which causes contractions to expel the pregnancy. For more information on medical abortion, visit genbiopro.com or scan here.



WE SUPPORT YOUR PREGNANCY DECISION

At GenBioPro, we believe that you are the expert on your own life. You are the only one with enough information to determine your future and make personal health care decisions best for you and your family. If medical abortion is right for you, we hope the recent changes to the prescribing guidelines for mifepristone will help you to access care.

New for You in 2023: Increased Access, Flexibility, and Convenience

UPDATED REGULATIONS

Mifepristone is regulated by the US Food & Drug Administration or FDA. Good news for you: in 2023 the FDA made changes to how medical abortion pills can be prescribed and dispensed. Now there are more ways for you to receive the care you need.

In-Person or Telemedicine Care: You Choose

In addition to in-person care, now telemedicine care is available in states where it is permitted. This gives you the flexibility to see your health care provider in person or remotely. You choose based on what is best for you.

More Ways to Get Your Prescription

Certified prescribers can now choose how to dispense generic mifepristone tablets, 200 mg — directly to you or by sending a prescription to a certified

pharmacy to be filled. Depending on the way your health care provider works, you may be able to receive generic mifepristone tablets, 200 mg, in person, pick up your prescription in a pharmacy, or a pharmacy may mail your prescription to you. This gives you more options for care. Ask your health care provider about the ways you can get your generic mifepristone tablets, 200 mg.

What do the changes to regulations for mifepristone mean to you? Increased access, more choice, and greater convenience.



HOW DO YOU GET GENERIC MIFEPRISTONE TABLETS, 200 MG?

It starts with a visit — in-person or by telemedicine, if permitted in the state where you are seeking care — to a health care provider. Generic mifepristone tablets, 200 mg, are not available over the counter. They are only available by prescription from health care providers who are GenBioPro certified prescribers. Before your health care provider prescribes mifepristone, you will be asked to review and sign an FDA-required Patient Agreement and Medication Guide. The Guide includes detailed instructions on how to take the pills, what to expect, potential risks of treatment, and contact information in case you experience any problems.

Ask your trusted health care provider about generic mifepristone tablets, 200 mg, or find an abortion care service near you.

FIND CARE WITH THESE TRUSTED RESOURCES

Planned Parenthood

Call 1-800-230-PLAN
[plannedparenthood.org](https://www.plannedparenthood.org)

National Abortion Federation

Call 1-800-772-9100
[prochoice.org](https://www.prochoice.org)

Abortion Care Network

Call 1-202-419-1444
[abortiocarenetwork.org](https://www.abortiocarenetwork.org)

Abortion Clinics Online

Call 1-800-455-8130
[abortionclinics.com](https://www.abortionclinics.com)

I Need an A

[ineedana.com](https://www.ineedana.com)

Abortion Finder

[abortionfinder.org](https://www.abortionfinder.org)

Here for You

GenBioPro is an innovative pharmaceutical company dedicated to putting reproductive health care within reach. Our company was founded on the belief that all people — regardless of income, gender, race, or geography — have a right to reproductive health care, including access to medical abortion. GenBioPro promotes personal dignity, access to medical evidence, self-determination, and high-quality health care for all. We are excited about the recent updates to the REMS guidelines for prescribing and dispensing generic mifepristone, 200 mg, and the advantages these changes bring to people like you.

Please see the Important Safety Information on page 3 and click here to read the [Medication Guide](#).

YOUR SOURCE FOR FDA-APPROVED GENERIC MIFEPRISTONE TABLETS, 200 MG



GenBioPro.com
Phone: 1-855-MIFE-INFO
Email: info@genbiopro.com

MIFEPRISTONE TABLETS, 200 MG

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT MIFEPRISTONE TABLETS, 200MG?

What symptoms should I be concerned with?

Although cramping and bleeding are an expected part of ending a pregnancy, rarely, serious and potentially life-threatening bleeding, infections, or other problems can occur following a miscarriage, surgical abortion, medical abortion, or childbirth. Seeking medical attention as soon as possible is needed in these circumstances. Serious infection has resulted in death in a very small number of cases. There is no information that use of Mifepristone tablets, 200 mg, and misoprostol caused these deaths. If you have any questions, concerns, or problems, or if you are worried about any side effects or symptoms, you should contact your healthcare provider.

Be sure to contact your health care provider promptly if you have any of the following:

Heavy Bleeding. Contact your healthcare provider right away if you bleed enough to soak through two thick full-size sanitary pads per hour for two consecutive hours or if you are concerned about heavy bleeding. In about 1 out of 100 women, bleeding can be so heavy that it requires a surgical procedure (surgical aspiration or D&C).

Abdominal Pain or "Feeling Sick." If you have abdominal pain or discomfort, or you are "feeling sick," including weakness, nausea, vomiting, or diarrhea, with or without fever, more than 24 hours after taking misoprostol, you should contact your healthcare provider without delay. These symptoms may be a sign of a serious infection or another problem (including an ectopic pregnancy, a pregnancy outside the womb).

Fever. In the days after treatment, if you have a fever of 100.4°F or higher that lasts for more than 4 hours, you should contact your healthcare provider right away. Fever may be a symptom of a serious infection or another problem.

If you cannot reach your healthcare provider, go to the nearest hospital emergency room.

What to do if you are still pregnant after mifepristone tablets, 200mg, with misoprostol treatment. If you are still pregnant, your healthcare provider will talk with you about a surgical procedure to end your pregnancy. In many cases, this surgical procedure can be done in the office/clinic. The chance of birth defects if the pregnancy is not ended is unknown.

Talk with your health care provider. Before you take Mifepristone tablets, 200 mg, you should read this [Medication Guide](#) and you and your healthcare provider should discuss the benefits and risks of your using Mifepristone tablets, 200 mg.

What is mifepristone tablets, 200mg?

Mifepristone tablets, 200mg, is used in a regimen with another prescription medicine called misoprostol to end an early pregnancy. Early pregnancy means it is 70 days (10 weeks) or less since your last menstrual period began. Mifepristone tablets, 200 mg is not approved for ending pregnancies that are further along. Mifepristone tablets, 200 mg blocks a hormone needed for your pregnancy to continue. When you use Mifepristone tablets,

200 mg on Day 1, you also need to take another medicine called misoprostol 24 to 48 hours after you take Mifepristone tablets, 200 mg to cause the pregnancy to be passed from your uterus.

The pregnancy is likely to be passed from your uterus within 2 to 24 hours after taking Mifepristone tablets, 200 mg and misoprostol. When the pregnancy is passed from the uterus, you will have bleeding and cramping that will likely be heavier than your usual period. About 2 to 7 out of 100 women taking Mifepristone tablets, 200 mg will need a surgical procedure because the pregnancy did not completely pass from the uterus or to stop bleeding.

Who should not take mifepristone tablets, 200mg?

Some patients should not take Mifepristone tablets, 200 mg. Do not take Mifepristone tablets, 200 mg if you:

- Have a pregnancy that is more than 70 days (10 weeks). Your healthcare provider may do a clinical examination, an ultrasound examination, or other testing to determine how far along you are in pregnancy.
- Are using an IUD (intrauterine device or system). It must be taken out before you take Mifepristone tablets, 200mg.
- Have been told by your health care provider you have a pregnancy outside the uterus (ectopic pregnancy).
- Have problems with your adrenal glands (chronic adrenal failure).
- Take medicine to thin your blood.
- Have a bleeding problem.
- Have porphyria.
- Take certain steroid medicines.
- Are allergic to mifepristone, misoprostol or medicines that contain misoprostol, such as Cytotec or Arthrotec.

Ask your health care provider if you are not sure about all your medical conditions before taking this medicine to find out if you can take Mifepristone tablets, 200mg.

What should I tell my health care provider before taking mifepristone tablets, 200mg?

Before you take Mifepristone tablets, 200mg, tell your health care provider if you:

- Cannot follow-up within approximately 7 to 14 days of your first visit.
- Are breastfeeding. Mifepristone tablets, 200mg, can pass into your breast milk. The effect of the Mifepristone tablets, 200mg, and misoprostol regimen on the breastfed infant or on milk production is unknown.
- Are taking medicines, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Mifepristone tablets, 200mg, and certain other medicines may affect each other if they are used together. This can cause side effects.

How should I take mifepristone tablets, 200mg?

Mifepristone tablets, 200 mg will be given to you by a healthcare provider or pharmacy.

You and your healthcare provider will plan the most appropriate

location for you to take the misoprostol, because it may cause bleeding, cramps, nausea, diarrhea, and other symptoms that usually begin within 2 to 24 hours after taking it.

Most women will pass the pregnancy within 2 to 24 hours after taking the misoprostol tablets.

Follow the instruction below on how to take Mifepristone tablets, 200mg, and misoprostol: Mifepristone tablets, 200mg, (1 tablet) orally + misoprostol (4 tablets) buccally

Day 1:

- Take 1 Mifepristone 200mg tablet by mouth.

24 to 48 hours after taking Mifepristone tablets, 200mg:

- Take 4 misoprostol tablets by placing 2 tablets in each cheek pouch (the area between your teeth and cheek—see Figure A) for 30 minutes and then swallow anything left over with a drink of water or another liquid.
- The medicines may not work as well if you take misoprostol sooner than 24 hours after Mifepristone tablets, 200 mg or later than 48 hours after Mifepristone tablets, 200 mg.
- Misoprostol often causes cramps, nausea, diarrhea, and other symptoms. Your healthcare provider may send you home with medicines for these symptoms.

Follow-up Assessment at Day 7 to 14:

- This follow-up assessment is very important. You must follow up with your healthcare provider about 7 to 14 days after you have taken Mifepristone tablets, 200 mg to be sure you are well, that you have had bleeding, and the pregnancy has passed from your uterus.
- Your healthcare provider will assess whether your pregnancy has passed from your uterus. If your pregnancy continues, the chance that there may be birth defects is unknown. If you are still pregnant, your healthcare provider will talk with you about a surgical procedure to end your pregnancy.
- If your pregnancy has ended, but not yet completely passed from your uterus, your provider will talk with you about other choices you have, including waiting, taking another dose of misoprostol, or having a surgical procedure to empty your uterus.

When should I begin birth control?

You can become pregnant again right after your pregnancy ends. If you do not want to become pregnant again, start using birth control as soon as your pregnancy ends or before you start having sexual intercourse again.

What should I avoid while taking Mifepristone tablets, 200 mg and misoprostol?

Do not take any other prescription or over-the-counter medicines (including herbal medicines or supplements) at any time during the treatment period without first asking your healthcare provider about them because they may interfere with the treatment. Ask your healthcare provider about what medicines you can take for pain and other side effects.

What are the possible side effects of Mifepristone tablets, 200mg and misoprostol?

Mifepristone tablets, 200 mg may cause serious side effects. See “What is the most important information I should know about Mifepristone tablets, 200 mg?”

Cramping and bleeding. Cramping and vaginal bleeding are expected with this treatment. Usually, these symptoms mean that the treatment is working. But sometimes you can get cramping and bleeding and still be pregnant. This is why you must follow up with your healthcare provider approximately 7 to 14 days after taking Mifepristone tablets, 200 mg. See “How should I take Mifepristone tablets, 200 mg?” for more information on your follow-up assessment. If you are not already bleeding after taking Mifepristone tablets, 200 mg, you probably will begin to bleed once you take misoprostol, the medicine you take 24 to 48 hours after Mifepristone tablets, 200 mg. Bleeding or spotting can be expected for an average of 9 to 16 days and may last for up to 30 days. Your bleeding may be similar to, or greater than, a normal heavy period. You may see blood clots and tissue. This is an expected part of passing the pregnancy.

The most common side effects of Mifepristone tablets, 200 mg treatment include: nausea, weakness, fever/chills, vomiting, headache, diarrhea, and dizziness. Your provider will tell you how to manage any pain or other side effects. These are not all the possible side effects of Mifepristone tablets, 200 mg.

Call your health care provider for medical advice about any side effects that bother you or do not go away. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of mifepristone tablets, 200mg.

Medicines are sometimes prescribed for purposes other than those listed in a [Medication Guide](#). This [Medication Guide](#) summarizes the most important information about Mifepristone tablets, 200 mg. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider for information about Mifepristone tablets, 200 mg that is written for healthcare professionals.

For additional information about mifepristone, see the [Medication Guide](#) and [Full Prescribing Information](#).

YOUR SOURCE FOR FDA-APPROVED GENERIC MIFEPRISTONE TABLETS, 200 MG



GenBioPro

PUTTING ACCESS INTO PRACTICE

GenBioPro.com

Phone: 1-855-MIFE-INFO

Email: info@genbiopro.com